

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Mesothelin (Mesothelial Marker) (MSLN/2131), 0.2mg/mL, Clone: [MSLN/2131], Mouse, Monoclonal BOT-BNUB2131-500

Artikelname	Mesothelin (Mesothelial Marker) (MSLN/2131), 0.2mg/mL, Clone: [MSLN/2131], Mouse, Monoclonal
Artikelnummer	BOT-BNUB2131-500
Hersteller Artikelnummer	BNUB2131-500
Alternativnummer	BOT-BNUB2131-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human, Mouse, Rat
Immunogen	Recombinant fragment (around aa 273-407) of human Mesothelin (MSLN) protein (exact sequence is proprietary)
Produktbeschreibung	Mesothelin is a 40 kDa glycosyl-phosphatidylinositol-anchored glycoprotein cleaved from a 69 kDa precursor protein. Mesothelin immunoreactivity is high in cancers of the ovary (serous papillary, endometrioid and undifferentiated) and pancreas, with I...
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[MSLN/2131]

Molekulargewicht	40 kDa (Mature form), 69 kDa (Precursor)
UniProt	Q13421
Puffer	PBS, 0.05% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user