

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

PU.1 (SPI-1) (B-Cell Marker) (PU1/2146), 0.2mg/mL, Clone: [PU1/2146], Mouse, Monoclonal BOT-BNUB2146-100

Artikelname	PU.1 (SPI-1) (B-Cell Marker) (PU1/2146), 0.2mg/mL, Clone: [PU1/2146], Mouse, Monoclonal
Artikelnummer	BOT-BNUB2146-100
Hersteller Artikelnummer	BNUB2146-100
Alternativnummer	BOT-BNUB2146-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC, IF, IHC, WB
Spezies Reaktivität	Human
Immunogen	Recombinant fragment (around aa 16-170) of human PU.1 protein (Exact sequence is proprietary)
Produktbeschreibung	PU.1 is a member of the ETS family of transcription factors and is important for normal B-cell development. It is expressed in the myeloid lineage and in immature as well as mature B-lymphocytes, with the exception of plasma cells. PU.1 is expressed ...
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[PU1/2146]

Molekulargewicht	40 kDa
UniProt	P17947
Puffer	PBS, 0.05% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Western Blot 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user