

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

MUC1 / CA15-3 / EMA / CD227 (Epithelial Marker) (MUC1/2278R), 0.2mg/mL, Clone: [MUC1/2278R], Rabbit, Monoclonal BOT-BNUB2278-100

Artikelname	MUC1 / CA15-3 / EMA / CD227 (Epithelial Marker) (MUC1/2278R), 0.2mg/mL, Clone: [MUC1/2278R], Rabbit, Monoclonal
Artikelnummer	BOT-BNUB2278-100
Hersteller Artikelnummer	BNUB2278-100
Alternativnummer	BOT-BNUB2278-100-100UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant full-length human MUC1 protein
Produktbeschreibung	This MAb reacts with MUC1, a large transmembrane glycoprotein expressed on the ductal surface of normal glandular epithelia. It is used as tracer agent in CA15.3 assays. The extracellular domain of MUC1 largely consists of a highly conserved, O-glyco...
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[MUC1/2278R]
Molekulargewicht	265-400 kDa

UniProt	P15941
Puffer	PBS, 0.05% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user