

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

p53(BP53-12), 1mg/mL, IgG2a, Clone: [BP53-12], Mouse, Monoclonal BOT-BNUM0013-50

Artikelname	p53(BP53-12), 1mg/mL, IgG2a, Clone: [BP53-12], Mouse, Monoclonal
Artikelnummer	BOT-BNUM0013-50
Hersteller Artikelnummer	BNUM0013-50
Alternativnummer	BOT-BNUM0013-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Canine, Gallus, Hamster, Human, Monkey
Immunogen	Recombinant human wild-type p53 protein
Produktbeschreibung	This MAb reacts with an N-terminal epitope (aa 16-25) of both wild type and mutated p53. Mutation and/or allelic loss of p53 is one of the causes of a variety of mesenchymal and epithelial tumors. If it occurs in the germ line, such tumors run in fam...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[BP53-12]
Molekulargewicht	53 kDa
Isotyp	IgG2a

UniProt	P04637
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Does not react with mouse or rat, others not known Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user