

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Granulocyte Marker(BM-2), 1mg/mL, IgG1, Clone: [BM-2], Mouse, Monoclonal BOT-BNUM0062-50

Artikelname	Granulocyte Marker(BM-2), 1mg/mL, IgG1, Clone: [BM-2], Mouse, Monoclonal
Artikelnummer	BOT-BNUM0062-50
Hersteller Artikelnummer	BNUM0062-50
Alternativnummer	BOT-BNUM0062-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Nuclei from pokeweed mitogen stimulated human peripheral blood lymphocytes
Produktbeschreibung	Recognizes 183 kDa protein with DNA-binding characteristics, which is identified as a myeloid specific antigen. It reacts with myeloid precursor cells and granulocytes in bone marrow. Its antigen appears to be restricted to M2 and M3 acute myelogenous...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[BM-2]

Molekulargewicht	183 kDa
Isotyp	IgG1
UniProt	Not Known
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user