

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Human Lambda Light Chain(LcN-2 + ICO-106), 1mg/mL, Clone: [LcN-2 ICO-106], Mouse, Monoclonal BOT-BNUM0735-50

Artikelname	Human Lambda Light Chain(LcN-2 + ICO-106), 1mg/mL, Clone: [LcN-2 ICO-106], Mouse, Monoclonal
Artikelnummer	BOT-BNUM0735-50
Hersteller Artikelnummer	BNUM0735-50
Alternativnummer	BOT-BNUM0735-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	WB
Spezies Reaktivität	Human
Immunogen	Purified human IgG (LcN-2 and ICO-106)
Produktbeschreibung	This MAb is specific to lambda light chain of immunoglobulin and shows no cross-reaction with lambda light chain or any of the five heavy chains. In mammals, the two light chains in an antibody are always identical, with only one type of light chain,...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[LcN-2 ICO-106]
Molekulargewicht	~22.5 kDa

UniProt	P01701
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1.0 ug/mL Optimal dilution for a specific application should be determined by user