

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Thomsen-Friedenreich Antigen / CD176 (Pan Carcinoma Marker)(A78-G/A7), 1mg/mL, Clone: [A78-G/A7], Mouse, Monoclonal BOT-BNUM0936-50

Artikelname	Thomsen-Friedenreich Antigen / CD176 (Pan Carcinoma Marker)(A78-G/A7), 1mg/mL, Clone: [A78-G/A7], Mouse, Monoclonal
Artikelnummer	BOT-BNUM0936-50
Hersteller Artikelnummer	BNUM0936-50
Alternativnummer	BOT-BNUM0936-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human, Mouse, Rat
Immunogen	Neuraminidase-treated human red blood cells
Produktbeschreibung	This antibody recognizes a disaccharide epitope, Gal 1-3GalNAc, of Thomsen-Friedenreich (TF) antigen. It is specific for both anomeric forms of the disaccharide (TF and TF , including related structures on the glycolipid) and shows no cross-reactivit...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[A78-G/A7]
Molekulargewicht	Multiple

UniProt	Not Applicable
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Immunohistochemistry (formalin-fixed): 0.5-1.0 ug/mL for 30 minutes at RT Immunofluorescence: 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user