

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Thymidine Phosphorylase / PD-ECGF (Angiogenesis Marker) (P-GF.44C), 1mg/mL, IgG1, Clone: [P-GF.44C], Mouse, Monoclonal BOT-BNUM1244-50

Artikelname	Thymidine Phosphorylase / PD-ECGF (Angiogenesis Marker) (P-GF.44C), 1mg/mL, IgG1, Clone: [P-GF.44C], Mouse, Monoclonal
Artikelnummer	BOT-BNUM1244-50
Hersteller Artikelnummer	BNUM1244-50
Alternativnummer	BOT-BNUM1244-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Human, Mouse, Rat
Immunogen	Recombinant full-length human Thymidine Phosphorylase (TP / PD-ECGF) protein
Produktbeschreibung	This antibody recognizes a protein (amino acid 482) of 55 kDa (in vivo 110 kDa homodimer), identified as platelet-derived endothelial growth factor (PD-ECGF), same as thymidine phosphorylase (TP) or gliostatin. In the presence of inorganic orthophosp...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[P-GF.44C]

Molekulargewicht	55 kDa
Isotyp	IgG1
UniProt	P19971
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin) 1-2 ug/mL Western blotting 0.5-1 ug/mL Immunoprecipitation 0.5-1 ug/500ug protein lysate Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Optimal dilution for a specific application should be determined by user