

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Napsin A (Lung Adenocarcinoma Marker)(NAPSA/1238 + NAPSA/1239), 1mg/mL, Clone: [NAPSA/1238 NAPSA/1239], Mouse, Monoclonal BOT-BNUM1254-50

Artikelname	Napsin A (Lung Adenocarcinoma Marker)(NAPSA/1238 + NAPSA/1239), 1mg/mL, Clone: [NAPSA/1238 NAPSA/1239], Mouse, Monoclonal
Artikelnummer	BOT-BNUM1254-50
Hersteller Artikelnummer	BNUM1254-50
Alternativnummer	BOT-BNUM1254-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Human
Immunogen	Recombinant human Napsin-A protein fragment (aa189-299) (exact sequence is proprietary)
Produktbeschreibung	Napsin is a pepsin-like aspartic proteinase connected with maturation of surfactant protein B. There are two closely related napsins, napsin A and napsin B. Napsin A is expressed as a single chain protein. Immunohistochemical studies revealed high exp...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[NAPSA/1238 NAPSA/1239]

Molekulargewicht	37 kDa
UniProt	O96009
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistochemistry (formalin-fixed): 1-2 ug/mL for 30 minutes at RT Western blot: 1-2 ug/mL for 60 minutes at RT Flow cytometry: 0.5-1 ug/million cells Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user