

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cytokeratin 18 (KRT18)(B23.1), 1mg/mL, Clone: [B23.1], Mouse, Monoclonal BOT-BNUM1263-50

Artikelname	Cytokeratin 18 (KRT18)(B23.1), 1mg/mL, Clone: [B23.1], Mouse, Monoclonal
Artikelnummer	BOT-BNUM1263-50
Hersteller Artikelnummer	BNUM1263-50
Alternativnummer	BOT-BNUM1263-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Bovine, Hamster, Human, Rat, Sheep
Immunogen	PMC-42 human breast carcinoma cells
Produktbeschreibung	This MAb reacts with a wide variety of simple epithelia. It does not react with stratified squamous epithelia. It reacts with epithelial tumors of the gastrointestinal tract, lung, breast, pancreas, ovary, and thyroid. Cytokeratin 18, which belongs t...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[B23.1]
Molekulargewicht	45 kDa

UniProt	P05783
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistochemistry (formalin-fixed): 0.5-1 ug/mL for 30 min at RT Flow cytometry: 0.5-1 ug/million cells Western Blot 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user