

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

c-Myc Oncoprotein(MYC275 + MYC909), 1mg/mL, Clone: [MYC275 MYC909], Mouse, Monoclonal BOT-BNUM1269-50

Artikelname	c-Myc Oncoprotein(MYC275 + MYC909), 1mg/mL, Clone: [MYC275 MYC909], Mouse, Monoclonal
Artikelnummer	BOT-BNUM1269-50
Hersteller Artikelnummer	BNUM1269-50
Alternativnummer	BOT-BNUM1269-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant human c-myc protein
Produktbeschreibung	This antibody recognizes a transcription factor of 64-67 kDa, identified as c-myc. This MAb shows no cross-reaction with v-myc. c-myc is involved in the control of cell proliferation and differentiation and is amplified and/or over-expressed in a var...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[MYC275 MYC909]
Molekulargewicht	62-64 kDa

UniProt	P01106
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistochemistry (formalin-fixed): 1-2 ug/mL for 30 minutes at RT Flow cytometry: 0.5-1 ug/million cells Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA pH 9.0 for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user