

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Ep-CAM / CD326 (Extracellular Domain) (Epithelial Marker) (EGP40/1373), 1mg/mL, IgG2b, Clone: [EGP40/1373], Mouse, Monoclonal BOT-BNUM1373-50

Artikelname	Ep-CAM / CD326 (Extracellular Domain) (Epithelial Marker) (EGP40/1373), 1mg/mL, IgG2b, Clone: [EGP40/1373], Mouse, Monoclonal
Artikelnummer	BOT-BNUM1373-50
Hersteller Artikelnummer	BNUM1373-50
Alternativnummer	BOT-BNUM1373-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IF, IHC, WB
Spezies Reaktivität	Human
Immunogen	Recombinant human EpCAM protein fragment from the extracellular domain (aa77-202) (exact sequence is proprietary)
Produktbeschreibung	EGP40 is a 40-43 kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep-CAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast majority...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[EGP40/1373]

Molekulargewicht	40-43 kDa
Isotyp	IgG2b
UniProt	P16422
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user