

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cytokeratin, Acidic (Type I or LMW) (Epithelial Marker) (KRTL/1377), 1mg/mL, IgG1, Clone: [KRTL/1377], Mouse, Monoclonal BOT-BNUM1377-50

Artikelname	Cytokeratin, Acidic (Type I or LMW) (Epithelial Marker) (KRTL/1377), 1mg/mL, IgG1, Clone: [KRTL/1377], Mouse, Monoclonal
Artikelnummer	BOT-BNUM1377-50
Hersteller Artikelnummer	BNUM1377-50
Alternativnummer	BOT-BNUM1377-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Spezies Reaktivität	Human
Immunogen	Recombinant human KRT77 protein fragment (exact sequence is proprietary)
Produktbeschreibung	This MAb recognizes the 56.5 kDa (CK10), 50 kDa (CK14), 50 kDa (CK15), 48 kDa (CK16), 40 kDa (CK19) keratins of the acidic (Type I or LMW) subfamily. Twenty human keratins are resolved with two-dimensional gel electrophoresis into acidic (pI 6.0) sub...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[KRTL/1377]
Molekulargewicht	56.5 kDa (CK10), 50 kDa (CK14), 50 kDa (CK15), 48 kDa (CK16), 40 kDa (CK19)

Isotyp	IgG1
UniProt	Q7Z794
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry