

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD44v4 (Marker of Tumor Metastasis) (CD44v4/1700R), 1mg/mL, Clone: [CD44v4/1700R], Rabbit, Monoclonal BOT-BNUM1700-50

Artikelname	CD44v4 (Marker of Tumor Metastasis) (CD44v4/1700R), 1mg/mL, Clone: [CD44v4/1700R], Rabbit, Monoclonal
Artikelnummer	BOT-BNUM1700-50
Hersteller Artikelnummer	BNUM1700-50
Alternativnummer	BOT-BNUM1700-50-50UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant fragment corresponding to the v4 domain of human CD44 (exact sequence is proprietary)
Produktbeschreibung	This antibody recognizes an epitope encoded by exon v4 on the variant portion of human CD44. The CD44 molecule belongs to a family of cellular adhesion molecules found on a wide range of normal and malignant cells in epithelial, mesothelial and hemop...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[CD44v4/1700R]

Molekulargewicht	80-95 kDa
UniProt	P16070
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user