

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD43 (T-Cell Marker) (SPN/1766R), 1mg/mL, Clone: [SPN/1766R], Rabbit, Monoclonal BOT-BNUM1766-50

Artikelname	CD43 (T-Cell Marker) (SPN/1766R), 1mg/mL, Clone: [SPN/1766R], Rabbit, Monoclonal
Artikelnummer	BOT-BNUM1766-50
Hersteller Artikelnummer	BNUM1766-50
Alternativnummer	BOT-BNUM1766-50-50UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	FC, IF, IHC, WB
Spezies Reaktivität	Human
Immunogen	Recombinant full-length human SPN protein
Produktbeschreibung	This antibody recognizes a cell surface glycoprotein of 95/115/135 kDa (depending upon the extent of glycosylation), identified as CD43. 70-90% of T-cell lymphomas and from 22-37% of B-cell lymphomas express CD43. No reactivity has been observed with...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[SPN/1766R]
Molekulargewicht	115 95 or 135 kDa

UniProt	P16150
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user