

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD22 / BL-CAM (BLCAM/1795), 1mg/mL, Clone: [BLCAM/1795], Mouse, Monoclonal BOT-BNUM1795-50

Artikelname	CD22 / BL-CAM (BLCAM/1795), 1mg/mL, Clone: [BLCAM/1795], Mouse, Monoclonal
Artikelnummer	BOT-BNUM1795-50
Hersteller Artikelnummer	BNUM1795-50
Alternativnummer	BOT-BNUM1795-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC, IF, IHC, WB
Spezies Reaktivität	Human
Immunogen	Recombinant human BL-CAM/CD22 protein (aa52-178) (exact sequence is proprietary)
Produktbeschreibung	This antibody recognizes a protein of 130-140 kDa, identified as CD22 (also known as BL-CAM). CD22 expression is restricted to normal and neoplastic B cells and is absent from other haemopoietic cell types. In B-cell ontogeny, CD22 is first expressed...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[BLCAM/1795]

Molekulargewicht	130-140 kDa
UniProt	P20273
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody ELISA: 2-4 ug/mL for coating order Ab without BSA Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin) 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris buffer with 1 mM EDTA pH 9.0 for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user