

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Human Kappa Light Chain / IGKC (B-Cell Marker) (rKLC709), 1mg/mL, Clone: [rKLC709], Mouse, Monoclonal BOT-BNUM2050-50

Artikelname	Human Kappa Light Chain / IGKC (B-Cell Marker) (rKLC709), 1mg/mL, Clone: [rKLC709], Mouse, Monoclonal
Artikelnummer	BOT-BNUM2050-50
Hersteller Artikelnummer	BNUM2050-50
Alternativnummer	BOT-BNUM2050-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant full-length human Ig kappa chain
Produktbeschreibung	This MAb is specific to kappa light chain of immunoglobulin and shows no cross-reaction with lambda light chain or any of the five heavy chains. It recognizes human Ig kappa light chains of both secreted and cell surface immunoglobulin. It detects al...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[rKLC709]
Molekulargewicht	~22.5 kDa

UniProt	P01601
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Flow cytometry: 0.5-1 ug/million cells Immunohistology (formalin): 0.5-1.0 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user