

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

PU.1 (SPI-1) (B-Cell Marker) (PU1/2146), 1mg/mL, Clone: [PU1/2146], Mouse, Monoclonal BOT-BNUM2146-50

Artikelname	PU.1 (SPI-1) (B-Cell Marker) (PU1/2146), 1mg/mL, Clone: [PU1/2146], Mouse, Monoclonal
Artikelnummer	BOT-BNUM2146-50
Hersteller Artikelnummer	BNUM2146-50
Alternativnummer	BOT-BNUM2146-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC, IF, IHC, WB
Spezies Reaktivität	Human
Immunogen	Recombinant fragment (around aa 16-170) of human PU.1 protein (Exact sequence is proprietary)
Produktbeschreibung	PU.1 is a member of the ETS family of transcription factors and is important for normal B-cell development. It is expressed in the myeloid lineage and in immature as well as mature B-lymphocytes, with the exception of plasma cells. PU.1 is expressed ...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[PU1/2146]

Molekulargewicht	40 kDa
UniProt	P17947
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Western Blot 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user