

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

NKX2.2 (Neuroendocrine & Ewing s Sarcoma Marker) (rNX2/1523), 1mg/mL, IgG1, Clone: [rNX2/1523], Mouse, Monoclonal BOT-BNUM2309-50

Artikelname	NKX2.2 (Neuroendocrine & Ewing s Sarcoma Marker) (rNX2/1523), 1mg/mL, IgG1, Clone: [rNX2/1523], Mouse, Monoclonal
Artikelnummer	BOT-BNUM2309-50
Hersteller Artikelnummer	BNUM2309-50
Alternativnummer	BOT-BNUM2309-50-50UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Human recombinant NKX2.2 protein fragment (around aa1-119) (exact sequence is proprietary)
Produktbeschreibung	Expression of NKX2.2 has been found in neuroendocrine tumors of the gut, making it a potential marker for the study of gastrointestinal neuroendocrine tumors. More recently, NKX2.2 protein was identified as a target of EWS-FLI-1, the fusion protein s...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[rNX2/1523]

Molekulargewicht	40-50 kDa
Isotyp	IgG1
UniProt	O95096
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user