

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

B7-H4 (Immuno-Inhibitory Protein) (B7H4/2652R), 1mg/mL, Clone: [B7H4/2652R], Rabbit, Monoclonal BOT-BNUM2652-50

Artikelname	B7-H4 (Immuno-Inhibitory Protein) (B7H4/2652R), 1mg/mL, Clone: [B7H4/2652R], Rabbit, Monoclonal
Artikelnummer	BOT-BNUM2652-50
Hersteller Artikelnummer	BNUM2652-50
Alternativnummer	BOT-BNUM2652-50-50UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	FC, IF, IHC
Spezies Reaktivität	Human
Immunogen	Recombinant human B7-H4 protein fragment (exact sequence is proprietary)
Produktbeschreibung	B7-H4 (B7-homolog 4), belongs to the B7 immunoglobulin superfamily of ligand-lymphocyte interacting proteins. Expressed primarily on the membrane of lymphoid cells, B7-H4 is an immuno-inhibitory protein that interacts with receptors on the surface of...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[B7H4/2652R]

Molekulargewicht	35 kDa
UniProt	Q7Z7D3
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody ELISA: 2-4 ug/m for coating order Ab without BSA Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user