

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Adipophilin / Perilipin-2 (Marker of Lipid Accumulation) (ADFP/2755R), 1mg/mL, Clone: [ADFP/2755R], Rabbit, Monoclonal BOT-BNUM2755-50

Artikelname	Adipophilin / Perilipin-2 (Marker of Lipid Accumulation) (ADFP/2755R), 1mg/mL, Clone: [ADFP/2755R], Rabbit, Monoclonal
Artikelnummer	BOT-BNUM2755-50
Hersteller Artikelnummer	BNUM2755-50
Alternativnummer	BOT-BNUM2755-50-50UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Human
Immunogen	Recombinant fragment (around aa 249-376) of human Adipophilin (ADFP) protein (exact sequence is proprietary)
Produktbeschreibung	Perilipin-2, also known as Adipophilin, belongs to the perilipin family, members of which coat intracellular lipid storage droplets. This 48 kDa protein is associated with the lipid globule surface membrane material, and maybe involved in development...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[ADFP/2755R]

Molekulargewicht	48 kDa
UniProt	Q99541
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Western blotting 1-2 ug/mL ELISA For coating, order Ab without BSA Optimal dilution for a specific application should be determined by user