

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

BCL10 (MALT-Lymphoma Marker) (BL10/2988R), 1mg/mL, Clone: [BL10/2988R], Rabbit, Monoclonal BOT-BNUM2988-50

Artikelname	BCL10 (MALT-Lymphoma Marker) (BL10/2988R), 1mg/mL, Clone: [BL10/2988R], Rabbit, Monoclonal
Artikelnummer	BOT-BNUM2988-50
Hersteller Artikelnummer	BNUM2988-50
Alternativnummer	BOT-BNUM2988-50-50UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	FC, IF, IHC
Spezies Reaktivität	Human
Immunogen	Recombinant human BCL10 protein fragment (around aa122-168) (exact sequence is proprietary)
Produktbeschreibung	B-cell lymphoma/leukemia 10 (BCL10) is a signaling protein that promotes apoptosis, pro-caspase-9 maturation, and activation of NF-kappa-B via NIK and IKK. BCL10 contains an N-terminal caspase recruitment domain (CARD), which is found in a number of ...
Klonalität	Monoclonal
Konzentration	1 mg/mL
Klon-Bezeichnung	[BL10/2988R]

Molekulargewicht	33 kDa
UniProt	O95999
Puffer	PBS, no BSA, no azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user