

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Donkey F(ab')<sub>2</sub> anti-Human IgG (H+L)-HRPO, MinX Bo,Ck,Go,Gp,Hs,Ho,Ms,Rb,Rt,Sh JIM-709-036-149**

|                                     |                                                                                             |
|-------------------------------------|---------------------------------------------------------------------------------------------|
| Artikelname                         | Donkey F(ab') <sub>2</sub> anti-Human IgG (H+L)-HRPO, MinX<br>Bo,Ck,Go,Gp,Hs,Ho,Ms,Rb,Rt,Sh |
| Artikelnummer                       | JIM-709-036-149                                                                             |
| Hersteller Artikelnummer            | 709-036-149                                                                                 |
| Alternativnummer                    | JIM-709-036-149                                                                             |
| Hersteller                          | Jackson ImmunoResearch                                                                      |
| Wirt                                | Donkey                                                                                      |
| Kategorie                           | Antikörper                                                                                  |
| Spezies Reaktivität                 | Human                                                                                       |
| Konjugation                         | HRPO                                                                                        |
| Format                              | F(ab') <sub>2</sub>                                                                         |
| Spezifität                          | IgG (H+L)                                                                                   |
| Minimale Kreuzreaktivität<br>(MinX) | Bovine,Gallus,Goat,Guinea pig,Hamster<br>(syrian),Equine,Mouse,Rabbit,Rat,Sheep             |
| Klonalität                          | Polyclonal                                                                                  |
| Isotyp                              | F(ab') <sub>2</sub> Fragment                                                                |

|                        |                                                                                                                                                                                                                                                                             |
|------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Puffer                 | <p>Buffer: 0.01M Sodium Phosphate, 0.25M NaCl, pH 7.6. Stabilizer: 15 mg/ml Bovine Serum Albumin (IgG-Free, Protease-Free).</p> <p>Preservative: None (Warning: Use of sodium azide as a preservative will substantially inhibit the enzyme activity of horseradish...</p>  |
| Reinheit               | <p>The antibody was purified from antisera by a combination of pepsin digestion and immunoaffinity chromatography using antigens coupled to agarose beads. Fc fragments and whole IgG molecules have been removed.</p>                                                      |
| Formulierung           | Freeze-dried solid                                                                                                                                                                                                                                                          |
| Lagerung               | <p>Storage and Rehydration: Store freeze-dried solid at 2-8°C.</p> <p>Rehydrate with the indicated volume of dH2O and centrifuge if not clear. Prepare working dilution on day of use. Product is stable for about 6 weeks at 2-8°C as an undiluted liquid. Extended...</p> |
| Antibody Type          | Secondary Antibody                                                                                                                                                                                                                                                          |
| Application Verdünnung | <p>1:500 - 1:5,000 for immunohisto/cytochemistry_x000D_ 1:5,000 - 1:100,000 for ELISA and Western blotting with chromogenic substrates_x000D_ 1:10,000 - 1:200,000 for Western blotting with ECL substrates_x000D_ Dilution factors are presented in the form...</p>        |