

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cytokeratin 7(KRT7/903), CF740 conjugate, 0.1mg/mL, IgG1, Clone: [KRT7/903], CF 740, Mouse, Monoclonal BOT-BNC740903-100

| | |
|--------------------------|---|
| Artikelname | Cytokeratin 7(KRT7/903), CF740 conjugate, 0.1mg/mL, IgG1, Clone: [KRT7/903], CF 740, Mouse, Monoclonal |
| Artikelnummer | BOT-BNC740903-100 |
| Hersteller Artikelnummer | BNC740903-100 |
| Alternativnummer | BOT-BNC740903-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC, IF, IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant full-length human KRT7 protein |
| Konjugation | CF 740 |
| Produktbeschreibung | Cytokeratin 7 is a basic cytokeratin, which is found in most glandular and transitional epithelia but not in the stratified squamous epithelia. Keratin 7 is expressed in the epithelial cells of ovary, lung, and breast but not of colon, prostate, or g... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [KRT7/903] |

| | |
|------------------------|--|
| Molekulargewicht | 54 kDa |
| Isotyp | IgG1 |
| UniProt | P08729 |
| Puffer | PBS, 0.1% rBSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 1-2 ug/mL Does not react with mouse, rat, ferret or rabbit, others not known Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris buffer with 1 mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user</p> |