

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Rat SIRT3 protein, His tag, Unconjugated GTX00061-PRO

Artikelname	Rat SIRT3 protein, His tag, Unconjugated
Artikelnummer	GTX00061-PRO
Hersteller Artikelnummer	GTX00061-pro
Alternativnummer	GTX00061-PRO-10
Hersteller	GeneTex
Kategorie	Proteine/Peptide
Applikation	FA
Spezies Reaktivität	Rat
Konjugation	Unconjugated
NCBI	293615
UniProt	C6ZII9
Puffer	Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300.
Expression System	E. coli
Formulierung	Lyophilized powder
Sequenz	N-terminal His-Tag, Pro127~Glu303

Anwendungsbeschreibung

Sirtuin 3 (SIRT3), the NAD-dependent deacetylase is a member of the mammalian sirtuin family of proteins. In human, sirtuins have a range of molecular functions and have emerged as important proteins in aging, stress resistance and metabolic regulation. It also can regulate epigenetic gene silencing and suppress recombination of rDNA in yeast. SIRT3 expression in white and brown adipose tissue. Besides, Isocitrate Dehydrogenase 2, mitochondrial (IDH2) has been identified as an interactor of SIRT3, thus a binding ELISA assay was conducted to detect the interaction of recombinant rat SIRT3 and recombinant rat IDH2. Briefly, SIRT3 were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100 µl were then transferred to IDH2-coated microtiter wells and incubated for 2h at 37°C. Wells were washed with PBST and incubated for 1h with anti-SIRT3 pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37°C. Finally, add 50 µl stop solution to the wells and read at 450nm immediately. The binding activity of SIRT3 and IDH2 was in a dose dependent manner.