

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Human TNFRSF14 protein, His tag, Unconjugated GTX00145-PRO

Artikelname	Human TNFRSF14 protein, His tag, Unconjugated
Artikelnummer	GTX00145-PRO
Hersteller Artikelnummer	GTX00145-pro
Alternativnummer	GTX00145-PRO-10
Hersteller	GeneTex
Kategorie	Proteine/Peptide
Applikation	FA
Spezies Reaktivität	Human
Konjugation	Unconjugated
NCBI	8764
UniProt	Q92956
Puffer	Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300.
Expression System	E. coli
Formulierung	Lyophilized powder
Sequenz	N-terminal His-Tag, Ser41~Ser209 (NP_003811.2)

Anwendungsbeschreibung	TNFRSF14 (Tumor necrosis factor receptor superfamily member 14) belongs to the tumor necrosis factor receptor superfamily. TNFRSF14 functions in signal transduction pathways that activate inflammatory and inhibitory T-cell immune response. It binds herpes simplex virus (HSV) viral envelope glycoprotein D (gD), mediating its entry into cells. A binding ELISA assay was conducted to detect the association of TNFRSF14 with TNFα. Briefly, TNFRSF14 were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100 μl TNFRSF14 were then transferred to TNFα-coated microtiter wells and incubated for 2h at 37C. Wells were washed with PBST and incubated for 1h with anti-TNFRSF14 pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37C. Finally, add 50 μl stop solution to the wells and read at 450nm immediately. The binding activity of TNFRSF14 and TNFα was in a dose dependent manner.
------------------------	---