

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Human MIF protein, His and GST tag, Unconjugated GTX00197-PRO

Artikelname	Human MIF protein, His and GST tag, Unconjugated
Artikelnummer	GTX00197-PRO
Hersteller Artikelnummer	GTX00197-pro
Alternativnummer	GTX00197-PRO-10
Hersteller	GeneTex
Kategorie	Proteine/Peptide
Applikation	FA
Spezies Reaktivität	Human
Konjugation	Unconjugated
NCBI	4282
UniProt	P14174
Puffer	Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300.
Expression System	E. coli
Formulierung	Lyophilized powder
Sequenz	N-terminal His and GST-Tag, Met3~Asn111 (NP_002406.1)

Anwendungsbeschreibung	Macrophage migration inhibitory factor (MIF), also known as glycosylation-inhibiting factor (GIF), L-dopachrome isomerase, or phenylpyruvate tautomerase is a protein classified as an inflammatory cytokine. MIF is an important regulator of innate immunity. It involved in cell-mediated immunity, immunoregulation, and inflammation. MIF plays a role in the regulation of macrophage function in host defense through the suppression of anti-inflammatory effects of glucocorticoids. This lymphokine and the JAB1 protein form a complex in the cytosol near the peripheral plasma membrane, which may indicate a role in integrin signaling pathways. Besides, Major Histocompatibility Complex Class II Invariant Chain (MHCDG) has been identified as an interactor of MIF, thus a binding ELISA assay was conducted to detect the interaction of recombinant human MIF and recombinant human MHCDG. Briefly, MIF were diluted serially in PBS, with 0.01% BSA (pH 7.4). Duplicate samples of 100 µl were then transferred to MHCDG-coated microtiter wells and incubated for 2h at 37C. Wells were washed with PBST and incubated for 1h with anti-MIF pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37C. Finally, add 50 µl stop solution to the wells and read at 450nm immediately. The binding activity of MIF and MHCDG was in a dose dependent manner.
------------------------	---