

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Human Her2 / ErbB2 protein, His and GST tag, Unconjugated GTX00232-PRO

Artikelname	Human Her2 / ErbB2 protein, His and GST tag, Unconjugated
Artikelnummer	GTX00232-PRO
Hersteller Artikelnummer	GTX00232-pro
Alternativnummer	GTX00232-PRO-10
Hersteller	GeneTex
Kategorie	Proteine/Peptide
Applikation	FA
Spezies Reaktivität	Human
Konjugation	Unconjugated
NCBI	2064
UniProt	P04626
Puffer	Reconstitute with 20mM Tris and 150mM NaCl to 0.1-1.0mg/ml. Do not vortex. Lyophilized from 20mM Tris, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose, ProClin 300.
Expression System	E. coli
Formulierung	Lyophilized powder
Sequenz	N-terminal His and GST-Tag, Phe376~Gly578 (NP_001005862.1)

Anwendungsbeschreibung

The epidermal growth factor receptor (EGFR, ErbB-1, HER1 in humans) is a transmembrane protein that is a receptor for members of the epidermal growth factor family (EGF family) of extracellular protein ligands. Epidermal growth factor receptor 2 (EGFR2) is a member of the epidermal growth factor (EGF) receptor family of receptor tyrosine kinases. The EGFR is essential for ductal development of the mammary glands, and agonists of the EGFR such as amphiregulin, TGF- α , and heregulin induce both ductal and lobuloalveolar development even in the absence of estrogen and progesterone. Besides, Actin Beta (ACTb) has been identified as an interactor of EGFR2, thus a binding ELISA assay was conducted to detect the interaction of recombinant human EGFR2 and recombinant human ACTb. Briefly, EGFR2 were diluted serially in PBS with 0.01% BSA (pH 7.4). Duplicate samples of 100 μ l were then transferred to ACTb-coated microtiter wells and incubated for 2h at 37C. Wells were washed with PBST and incubated for 1h with anti-EGFR2 pAb, then aspirated and washed 3 times. After incubation with HRP labelled secondary antibody, wells were aspirated and washed 3 times. With the addition of substrate solution, wells were incubated 15-25 minutes at 37C. Finally, add 50 μ l stop solution to the wells and read at 450nm immediately. The binding activity of EGFR2 and ACTb was in a dose dependent manner.