

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cytokeratin 7 Antibody Cocktail, Clone: [KRT7/760 + OV-TL12/30], Mouse, Monoclonal NSJ-V2660IHC-7ML

| | |
|--------------------------|---|
| Artikelname | Cytokeratin 7 Antibody Cocktail, Clone: [KRT7/760 + OV-TL12/30], Mouse, Monoclonal |
| Artikelnummer | NSJ-V2660IHC-7ML |
| Hersteller Artikelnummer | V2660IHC-7ML |
| Alternativnummer | NSJ-V2660IHC-7ML |
| Hersteller | NSJ Bioreagents |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC-P |
| Spezies Reaktivität | Human, Rat |
| Immunogen | Recombinant full-length human KRT7 protein (KRT7/760) and OTN 11 cells (OV-TL12/30) were used as the immunogen for the Cytokeratin 7 antibody cocktail. |
| Produktbeschreibung | It recognizes an intermediate filament protein (IFP) of 55kDa, which is identified as cytokeratin 7. This mAb is highly specific to cytokeratin 7 and shows no cross-reaction with other IFPs. Cytokeratin 7 is a basic cytokeratin, which is found in mos... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [KRT7/760 + OV-TL12/30] |
| UniProt | P08729 |

| | |
|------------------------|---|
| Reinheit | Protein G affinity chromatography |
| Formulierung | Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only* |
| Antibody Type | Primary Antibody |
| Application Verdünnung | The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |
| Anwendungsbeschreibung | Optimal dilution of the Cytokeratin 7 antibody cocktail should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min ² . The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |