

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cytokeratin 8 Antibody, Clone: [C-43], Mouse, Monoclonal NSJ-V2663IHC-7ML

| | |
|--------------------------|---|
| Artikelname | Cytokeratin 8 Antibody, Clone: [C-43], Mouse, Monoclonal |
| Artikelnummer | NSJ-V2663IHC-7ML |
| Hersteller Artikelnummer | V2663IHC-7ML |
| Alternativnummer | NSJ-V2663IHC-7ML |
| Hersteller | NSJ Bioreagents |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC-P |
| Spezies Reaktivität | Human |
| Immunogen | A cytoskeletal preparation from HeLa cells was used as the immunogen for the Cytokeratin 8 antibody. |
| Produktbeschreibung | Cytokeratin 8 (CK8) belongs to the type II (or B or basic) subfamily of high molecular weight cytokeratins and exists in combination with cytokeratin 18 (CK18). CK8 is primarily found in the non-squamous epithelia and is present in majority of adenoc... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [C-43] |
| UniProt | P05787 |
| Reinheit | Protein G affinity chromatography |

| | |
|------------------------|---|
| Formulierung | Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only* |
| Antibody Type | Primary Antibody |
| Application Verdünnung | The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |
| Anwendungsbeschreibung | Optimal dilution of the Cytokeratin 8 antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |