

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Cytokeratin 8 Antibody, Clone: [C-43], Mouse, Monoclonal NSJ-V2663SAF-100UG

Artikelname	Cytokeratin 8 Antibody, Clone: [C-43], Mouse, Monoclonal
Artikelnummer	NSJ-V2663SAF-100UG
Hersteller Artikelnummer	V2663SAF-100UG
Alternativnummer	NSJ-V2663SAF-100UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IF, IHC-P, WB
Spezies Reaktivität	Human
Immunogen	A cytoskeletal preparation from HeLa cells was used as the immunogen for the Cytokeratin 8 antibody.
Produktbeschreibung	Cytokeratin 8 (CK8) belongs to the type II (or B or basic) subfamily of high molecular weight cytokeratins and exists in combination with cytokeratin 18 (CK18). CK8 is primarily found in the non-squamous epithelia and is present in majority of adenoc...
Klonalität	Monoclonal
Klon-Bezeichnung	[C-43]
UniProt	<a href="#">P05787</a>
Reinheit	Protein G affinity chromatography

Formulierung	1 mg/ml in 1X PBS, BSA free, sodium azide free
Antibody Type	Primary Antibody
Application Verdünnung	Immunofluorescence: 1-2ug/ml,Western blot: 1-2ug/ml,Immunohistochemistry (FFPE): 0.5-1ug/ml for 30 min at RT
Anwendungsbeschreibung	Optimal dilution of the Cytokeratin 8 antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.