

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Keratin 8 Antibody / Cytokeratin 8 / CK8, Clone: [KRT8/803], Mouse, Monoclonal NSJ-V2664IHC-7ML**

Artikelname	Keratin 8 Antibody / Cytokeratin 8 / CK8, Clone: [KRT8/803], Mouse, Monoclonal
Artikelnummer	NSJ-V2664IHC-7ML
Hersteller Artikelnummer	V2664IHC-7ML
Alternativnummer	NSJ-V2664IHC-7ML
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Recombinant human KRT8 protein was used as the immunogen for the Keratin 8 antibody.
Produktbeschreibung	Cytokeratin 8 (CK8) belongs to the type II (or B or basic) subfamily of high molecular weight cytokeratins and exists in combination with cytokeratin 18 (CK18). CK8 is primarily found in the non-squamous epithelia and is present in majority of adenoc...
Klonalität	Monoclonal
Klon-Bezeichnung	[KRT8/803]
UniProt	<a href="#">P05787</a>

Reinheit	Protein G affinity chromatography
Formulierung	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Verdünnung	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Anwendungsbeschreibung	Optimal dilution of the Keratin 8 antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.