

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Keratin 14 Antibody, Clone: [KRT14/532], Mouse, Monoclonal NSJ-V2668-20UG**

|                          |   |
|--------------------------|---|
| Artikelname              | Keratin 14 Antibody, Clone: [KRT14/532], Mouse, Monoclonal  |
| Artikelnummer            | NSJ-V2668-20UG  |
| Hersteller Artikelnummer | V2668-20UG  |
| Alternativnummer         | NSJ-V2668-20UG  |
| Hersteller               | NSJ Bioreagents   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | IHC-P   |
| Spezies Reaktivität      | Human, Mouse, Rat   |
| Immunogen                | Recombinant full-length human KRT14 protein was used as the immunogen for the Keratin 14 antibody.  |
| Produktbeschreibung      | Cytokeratin 14 (CK14) belongs to the type I (or A or acidic) subfamily of low molecular weight keratins and exists in combination with keratin 5 (type II or B or basic). CK14 is found in basal cells of squamous epithelia, some glandular epithelia, my... |
| Klonalität               | Monoclonal  |
| Klon-Bezeichnung         | [KRT14/532]   |
| UniProt                  | <a href="#">P02533</a>  |
| Reinheit                 | Protein G affinity chromatography   |

|                        |  |
|------------------------|--|
| Formulierung           | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide   |
| Antibody Type          | Primary Antibody   |
| Application Verdünnung | Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT (1),Prediluted IHC only format: incubate for 30 min at RT (2)   |
| Anwendungsbeschreibung | Optimal dilution of the Keratin 14 antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |