

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cytokeratin 17 Antibody (KRT17), Clone: [KRT17/778], Mouse, Monoclonal NSJ-V2669-20UG

Artikelname	Cytokeratin 17 Antibody (KRT17), Clone: [KRT17/778], Mouse, Monoclonal
Artikelnummer	NSJ-V2669-20UG
Hersteller Artikelnummer	V2669-20UG
Alternativnummer	NSJ-V2669-20UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human, Rat
Immunogen	Recombinant full-length human KRT17 protein was used as the immunogen for the Cytokeratin 17 antibody.
Produktbeschreibung	Cytokeratin 17 (CK17) is normally expressed in the basal cells of complex epithelia but not in stratified or simple epithelia. Antibody to CK17 is an excellent tool to distinguish myoepithelial cells from luminal epithelium of various glands such as ...
Klonalität	Monoclonal
Klon-Bezeichnung	[KRT17/778]
UniProt	Q04695
Reinheit	Protein G affinity chromatography

Formulierung	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Verdünnung	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Anwendungsbeschreibung	Optimal dilution of the Cytokeratin 17 antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min ² . The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.