

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Keratin 18 Antibody, Clone: [KRT18/1190], Mouse, Monoclonal NSJ-V2670-20UG

| | |
|--------------------------|---|
| Artikelname | Keratin 18 Antibody, Clone: [KRT18/1190], Mouse, Monoclonal |
| Artikelnummer | NSJ-V2670-20UG |
| Hersteller Artikelnummer | V2670-20UG |
| Alternativnummer | NSJ-V2670-20UG |
| Hersteller | NSJ Bioreagents |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IF, IHC-P, WB |
| Spezies Reaktivität | Human, Rat |
| Immunogen | Recombinant human KRT18 protein was used as the immunogen for the Keratin 18 antibody. |
| Produktbeschreibung | This mAb reacts with a wide variety of simple epithelia. It does not react with stratified squamous epithelia. It reacts with epithelial tumors of the gastrointestinal tract, lung, breast, pancreas, ovary, and thyroid. Cytokeratin 18, which belongs t... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [KRT18/1190] |
| UniProt | P05783 |
| Reinheit | Protein G affinity chromatography |

| | |
|------------------------|--|
| Formulierung | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide |
| Antibody Type | Primary Antibody |
| Application Verdünnung | Immunofluorescence: 1-2ug/ml,Western blot: 1-2ug/ml,Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT |
| Anwendungsbeschreibung | Optimal dilution of the Keratin 18 antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |