

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CK18 Antibody / Cytokeratin 18, Clone: [KRT18/836], Mouse, Monoclonal NSJ-V2674-100UG

| | |
|--------------------------|---|
| Artikelname | CK18 Antibody / Cytokeratin 18, Clone: [KRT18/836], Mouse, Monoclonal |
| Artikelnummer | NSJ-V2674-100UG |
| Hersteller Artikelnummer | V2674-100UG |
| Alternativnummer | NSJ-V2674-100UG |
| Hersteller | NSJ Bioreagents |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IF, IHC-P, WB |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human KRT18 protein was used as the immunogen for the CK18 antibody. |
| Produktbeschreibung | This mAb reacts with a wide variety of simple epithelia. It does not react with stratified squamous epithelia. It reacts with epithelial tumors of the gastrointestinal tract, lung, breast, pancreas, ovary, and thyroid. Cytokeratin 18, which belongs t... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [KRT18/836] |
| UniProt | P05783 |
| Reinheit | Protein G affinity chromatography |

| | |
|------------------------|---|
| Formulierung | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide |
| Antibody Type | Primary Antibody |
| Application Verdünnung | Western blot: 1-2ug/ml,Immunofluorescence: 1-2ug/ml,Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT (1) |
| Anwendungsbeschreibung | Optimal dilution of the CK18 antibody should be determined by the researcher.1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |