

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cytokeratin 19 Antibody, Clone: [KRT19/799], Mouse, Monoclonal NSJ-V2678SAF-100UG

| | |
|--------------------------|---|
| Artikelname | Cytokeratin 19 Antibody, Clone: [KRT19/799], Mouse, Monoclonal |
| Artikelnummer | NSJ-V2678SAF-100UG |
| Hersteller Artikelnummer | V2678SAF-100UG |
| Alternativnummer | NSJ-V2678SAF-100UG |
| Hersteller | NSJ Bioreagents |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FACS, IF, IHC-P, WB |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human KRT19 protein was used as the immunogen for the Cytokeratin 19 antibody. |
| Produktbeschreibung | Recognizes a protein of 40kDa, identified as cytokeratin-19 (CK19), which is expressed in sweat gland, mammary gland ductal and secretory cells, bile ducts, gastrointestinal tract, bladder urothelium, oral epithelia, esophagus, and ectocervical epith... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [KRT19/799] |
| UniProt | P08727 |
| Reinheit | Protein G affinity chromatography |

| | |
|------------------------|---|
| Formulierung | 1 mg/ml in 1X PBS, BSA free, sodium azide free |
| Antibody Type | Primary Antibody |
| Application Verdünnung | Flow cytometry: 1-2ug/10 ⁶ cells, Immunofluorescence: 1-2ug/ml, Western blot: 1-2ug/ml, Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT |
| Anwendungsbeschreibung | Optimal dilution of the Cytokeratin 19 antibody should be determined by the researcher. 1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min. 2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |