

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CK19 Antibody / Cytokeratin 19, Clone: [KRT19/800], Mouse, Monoclonal NSJ-V2679-100UG

| | |
|--------------------------|---|
| Artikelname | CK19 Antibody / Cytokeratin 19, Clone: [KRT19/800], Mouse, Monoclonal |
| Artikelnummer | NSJ-V2679-100UG |
| Hersteller Artikelnummer | V2679-100UG |
| Alternativnummer | NSJ-V2679-100UG |
| Hersteller | NSJ Bioreagents |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IF, IHC-P, WB |
| Spezies Reaktivität | Human, Rat |
| Immunogen | Recombinant human Cytokeratin 19 protein was used as the immunogen for the CK19 antibody. |
| Produktbeschreibung | Recognizes a protein of 40kDa, identified as cytokeratin-19 (CK19), which is expressed in sweat gland, mammary gland ductal and secretory cells, bile ducts, gastrointestinal tract, bladder urothelium, oral epithelia, esophagus, and ectocervical epith... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [KRT19/800] |
| UniProt | P08727 |
| Reinheit | Protein G affinity chromatography |

| | |
|------------------------|--|
| Formulierung | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide |
| Antibody Type | Primary Antibody |
| Application Verdünnung | Western blot: 1-2ug/million cells, Immunofluorescence: 1-2ug/ml, Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT |
| Anwendungsbeschreibung | Optimal dilution of the CK19 antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |