

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CK19 Antibody / Cytokeratin 19, Clone: [KRT19/800], Mouse, Monoclonal NSJ-V2679IHC-7ML

Artikelname	CK19 Antibody / Cytokeratin 19, Clone: [KRT19/800], Mouse, Monoclonal
Artikelnummer	NSJ-V2679IHC-7ML
Hersteller Artikelnummer	V2679IHC-7ML
Alternativnummer	NSJ-V2679IHC-7ML
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human, Rat
Immunogen	Recombinant human Cytokeratin 19 protein was used as the immunogen for the CK19 antibody.
Produktbeschreibung	Recognizes a protein of 40kDa, identified as cytokeratin-19 (CK19), which is expressed in sweat gland, mammary gland ductal and secretory cells, bile ducts, gastrointestinal tract, bladder urothelium, oral epithelia, esophagus, and ectocervical epith...
Klonalität	Monoclonal
Klon-Bezeichnung	[KRT19/800]
UniProt	P08727
Reinheit	Protein G affinity chromatography

Formulierung	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Verdünnung	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Anwendungsbeschreibung	Optimal dilution of the CK19 antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.