

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

KRT19 Antibody / Cytokeratin 19, Clone: [Ks19.1], Mouse, Monoclonal NSJ-V2680-20UG

Artikelname	KRT19 Antibody / Cytokeratin 19, Clone: [Ks19.1], Mouse, Monoclonal
Artikelnummer	NSJ-V2680-20UG
Hersteller Artikelnummer	V2680-20UG
Alternativnummer	NSJ-V2680-20UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IF, IHC-P, WB
Spezies Reaktivität	Human
Immunogen	Human breast cancer MCF-7 cells were used as the immunogen for the KRT19 antibody.
Produktbeschreibung	This mAb reacts with the rod domain of human cytokeratin-19 (CK19), a polypeptide of 40kDa. Its epitope maps between amino acid 312-335. CK19 is expressed in sweat gland, mammary gland ductal and secretory cells, bile ducts, gastrointestinal tract, b...
Klonalität	Monoclonal
Klon-Bezeichnung	[Ks19.1]
UniProt	P08727
Reinheit	Protein G affinity chromatography

Formulierung	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Verdünnung	Western blot: 1-2ug/10 ⁶ cells, Immunofluorescence: 1-2ug/ml, Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Anwendungsbeschreibung	Optimal dilution of the KRT19 antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.