

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

KRT19 Antibody / Cytokeratin 19, Clone: [Ks19.1], Mouse, Monoclonal NSJ-V2680IHC-7ML

Artikelname	KRT19 Antibody / Cytokeratin 19, Clone: [Ks19.1], Mouse, Monoclonal
Artikelnummer	NSJ-V2680IHC-7ML
Hersteller Artikelnummer	V2680IHC-7ML
Alternativnummer	NSJ-V2680IHC-7ML
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Human breast cancer MCF-7 cells were used as the immunogen for the KRT19 antibody.
Produktbeschreibung	This mAb reacts with the rod domain of human cytokeratin-19 (CK19), a polypeptide of 40kDa. Its epitope maps between amino acid 312-335. CK19 is expressed in sweat gland, mammary gland ductal and secretory cells, bile ducts, gastrointestinal tract, b...
Klonalität	Monoclonal
Klon-Bezeichnung	[Ks19.1]
UniProt	P08727
Reinheit	Protein G affinity chromatography

Formulierung	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Verdünnung	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Anwendungsbeschreibung	Optimal dilution of the KRT19 antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.