

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Laminin gamma 1 Antibody / LAMC1 / LAMB2, Rat, Monoclonal NSJ-V2682-100UG

Artikelname	Laminin gamma 1 Antibody / LAMC1 / LAMB2, Rat, Monoclonal
Artikelnummer	NSJ-V2682-100UG
Hersteller Artikelnummer	V2682-100UG
Alternativnummer	NSJ-V2682-100UG
Hersteller	NSJ Bioreagents
Wirt	Rat
Kategorie	Antikörper
Applikation	FACS, IF, IHC-P
Spezies Reaktivität	Human, Mouse
Immunogen	A murine EHS laminin preparation was used as the immunogen for the Laminin gamma 1 antibody.
Produktbeschreibung	Laminins are large hetero-trimeric, non-collagenous glycoproteins composed of alpha, beta, and gamma chains. This mAb reacts with laminin B2/1 chain of ~210kDa and does not cross-react with other basement membrane components or fibronectin. Its speci...
Klonalität	Monoclonal
UniProt	P11047
Reinheit	Protein G affinity chromatography
Formulierung	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide

Antibody Type	Primary Antibody
Application Verdünnung	Flow cytometry: 0.5-1ug/10 ⁶ cells, Immunofluorescence: 0.5-1ug/ml, Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Anwendungsbeschreibung	Optimal dilution of the Laminin gamma 1 antibody should be determined by the researcher. 1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min. 2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.