

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### Laminin gamma 1 Antibody / LAMC1 / LAMB2, Rat, Monoclonal NSJ-V2682IHC-7ML

Artikelname	Laminin gamma 1 Antibody / LAMC1 / LAMB2, Rat, Monoclonal
Artikelnummer	NSJ-V2682IHC-7ML
Hersteller Artikelnummer	V2682IHC-7ML
Alternativnummer	NSJ-V2682IHC-7ML
Hersteller	NSJ Bioreagents
Wirt	Rat
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human, Mouse
Immunogen	A murine EHS laminin preparation was used as the immunogen for the Laminin gamma 1 antibody.
Produktbeschreibung	Laminins are large hetero-trimeric, non-collagenous glycoproteins composed of alpha, beta, and gamma chains. This mAb reacts with laminin B2/1 chain of ~210kDa and does not cross-react with other basement membrane components or fibronectin. Its speci...
Klonalität	Monoclonal
UniProt	<a href="#">P11047</a>
Reinheit	Protein G affinity chromatography
Formulierung	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*

Antibody Type	Primary Antibody
Application Verdünnung	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Anwendungsbeschreibung	Optimal dilution of the Laminin gamma 1 antibody should be determined by the researcher. <sup>1</sup> Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min <sup>2</sup> . The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.