

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Luteinizing Hormone beta Antibody, Clone: [SPM103], Mouse, Monoclonal NSJ-V2685-20UG

| | |
|--------------------------|---|
| Artikelname | Luteinizing Hormone beta Antibody, Clone: [SPM103], Mouse, Monoclonal |
| Artikelnummer | NSJ-V2685-20UG |
| Hersteller Artikelnummer | V2685-20UG |
| Alternativnummer | NSJ-V2685-20UG |
| Hersteller | NSJ Bioreagents |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC-P |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant beta sub-unit of human LH was used as the immunogen for the Luteinizing Hormone beta antibody. |
| Produktbeschreibung | Luteinizing hormone (LH) is a glycoprotein. Each monomeric unit is a sugar-like protein molecule, two of these make the full, functional protein. Its structure is similar to the other glycoproteins, follicle-stimulating hormone (FSH), thyroid-stimula... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [SPM103] |
| UniProt | P01229 |
| Reinheit | Protein G affinity chromatography |

| | |
|------------------------|--|
| Formulierung | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide |
| Antibody Type | Primary Antibody |
| Application Verdünnung | Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT |
| Anwendungsbeschreibung | Optimal dilution of the Luteinizing Hormone beta antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min |