

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

EpCAM Antibody, Clone: [EGP40/1120], Mouse, Monoclonal NSJ-V2687IHC-7ML

| | |
|--------------------------|---|
| Artikelname | EpCAM Antibody, Clone: [EGP40/1120], Mouse, Monoclonal |
| Artikelnummer | NSJ-V2687IHC-7ML |
| Hersteller Artikelnummer | V2687IHC-7ML |
| Alternativnummer | NSJ-V2687IHC-7ML |
| Hersteller | NSJ Bioreagents |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC-P |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant full-length human protein was used as the immunogen for the EpCAM antibody. |
| Produktbeschreibung | EGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep-CAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast major... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [EGP40/1120] |
| UniProt | P16422 |
| Reinheit | Protein G affinity chromatography |

| | |
|------------------------|---|
| Formulierung | Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only* |
| Antibody Type | Primary Antibody |
| Application Verdünnung | The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |
| Anwendungsbeschreibung | Optimal dilution of the EpCAM antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |