

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

EpCAM Antibody, Clone: [MOC-31], Mouse, Monoclonal NSJ-V2690-20UG

| | |
|--------------------------|---|
| Artikelname | EpCAM Antibody, Clone: [MOC-31], Mouse, Monoclonal |
| Artikelnummer | NSJ-V2690-20UG |
| Hersteller Artikelnummer | V2690-20UG |
| Alternativnummer | NSJ-V2690-20UG |
| Hersteller | NSJ Bioreagents |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC-P, WB |
| Spezies Reaktivität | Human |
| Immunogen | Neuraminidase treated GLS-1 human small cell lung carcinoma cells were used as the immunogen for the EpCAM antibody. |
| Produktbeschreibung | Binding epitope of this antibody is located in the first EGF-like repeat domain (EGF1) between amino acids 27-59 of Ep-CAM. EGP40 is a 40-43 kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelia... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [MOC-31] |
| UniProt | P16422 |
| Reinheit | Protein G affinity chromatography |

| | |
|------------------------|---|
| Formulierung | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide |
| Antibody Type | Primary Antibody |
| Application Verdünnung | Western blot: 1-2ug/ml, Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT |
| Anwendungsbeschreibung | <p>Optimal dilution of the EpCAM antibody should be determined by the researcher.</p> <ol style="list-style-type: none">1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.3. View the recombinant version of this recombinant-epcam-antibody-mouse-monoclonal-rmoc-31-v3464 EpCAM antibody. |