

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

TG Antibody / Thyroglobulin, Clone: [TGB24], Mouse, Monoclonal NSJ-V2883-100UG

| | |
|--------------------------|--|
| Artikelname | TG Antibody / Thyroglobulin, Clone: [TGB24], Mouse, Monoclonal |
| Artikelnummer | NSJ-V2883-100UG |
| Hersteller Artikelnummer | V2883-100UG |
| Alternativnummer | NSJ-V2883-100UG |
| Hersteller | NSJ Bioreagents |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | ELISA, IHC-P |
| Spezies Reaktivität | Human |
| Immunogen | Human thyroid follicular cells were used as the immunogen for the TG antibody. |
| Produktbeschreibung | Thyroglobulin is a 660 kDa dimeric pre-protein with multiple glycosylation sites. It is produced by and processed within the thyroid gland to produce the hormone thyroxine and triiodothyronine. Prior to forming dimers, thyroglobulin monomers undergo c... |
| Klonalität | Monoclonal |
| Klon-Bezeichnung | [TGB24] |
| UniProt | P01266 |
| Reinheit | Protein G affinity chromatography |

| | |
|------------------------|---|
| Formulierung | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide |
| Antibody Type | Primary Antibody |
| Application Verdünnung | ELISA: 2-4ug/ml (order BSA/azide-free format),Immunohistochemistry (FFPE): 0.5-1ug/ml for 30 min at RT |
| Anwendungsbeschreibung | Optimal dilution of the TG antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min. |