

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### TG Antibody / Thyroglobulin, Clone: [TGB24], Mouse, Monoclonal NSJ-V2883SAF-100UG

Artikelname	TG Antibody / Thyroglobulin, Clone: [TGB24], Mouse, Monoclonal
Artikelnummer	NSJ-V2883SAF-100UG
Hersteller Artikelnummer	V2883SAF-100UG
Alternativnummer	NSJ-V2883SAF-100UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	ELISA, IHC-P
Spezies Reaktivität	Human
Immunogen	Human thyroid follicular cells were used as the immunogen for the TG antibody.
Produktbeschreibung	Thyroglobulin is a 660 kDa dimeric pre-protein with multiple glycosylation sites. It is produced by and processed within the thyroid gland to produce the hormone thyroxine and triiodothyronine. Prior to forming dimers, thyroglobulin monomers undergo c...
Klonalität	Monoclonal
Klon-Bezeichnung	[TGB24]
UniProt	<a href="#">P01266</a>
Reinheit	Protein G affinity chromatography

Formulierung	1 mg/ml in 1X PBS, BSA free, sodium azide free
Antibody Type	Primary Antibody
Application Verdünnung	ELISA: 2-4ug/ml (order BSA/azide-free format),Immunohistochemistry (FFPE): 0.5-1ug/ml for 30 min at RT
Anwendungsbeschreibung	Optimal dilution of the TG antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.