

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### TGFa Antibody, Clone: [TG86], Mouse, Monoclonal NSJ-V2885-20UG

Artikelname	TGFa Antibody, Clone: [TG86], Mouse, Monoclonal
Artikelnummer	NSJ-V2885-20UG
Hersteller Artikelnummer	V2885-20UG
Alternativnummer	NSJ-V2885-20UG
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	FACS, IF, IHC-P
Spezies Reaktivität	Human
Immunogen	Amino acids 34-43 (PPVAAAVVSH) from the human protein were used as the immunogen for the TGFa antibody.
Produktbeschreibung	This antibody reacts with the TGF alpha and shows no cross-reaction with EGF and the neuropeptide synenkephalin. The staining with this mAb is completely blocked by the peptide used for raising this antibody. TGFa is a growth factor with 33% homology...
Klonalität	Monoclonal
Klon-Bezeichnung	[TG86]
UniProt	<a href="#">P01135</a>
Reinheit	Protein G affinity chromatography

Formulierung	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide
Antibody Type	Primary Antibody
Application Verdünnung	Flow cytometry: 0.5-1ug/10 <sup>6</sup> cells, Immunofluorescence: 1-2ug/ml, Immunohistochemistry (FFPE): 2-4ug/ml for 30 min at RT
Anwendungsbeschreibung	Optimal dilution of the TGFa antibody should be determined by the researcher. 1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min.