

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

TGM2 Antibody, Clone: [SPM592], Mouse, Monoclonal NSJ-V2891IHC-7ML

Artikelname	TGM2 Antibody, Clone: [SPM592], Mouse, Monoclonal
Artikelnummer	NSJ-V2891IHC-7ML
Hersteller Artikelnummer	V2891IHC-7ML
Alternativnummer	NSJ-V2891IHC-7ML
Hersteller	NSJ Bioreagents
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC-P
Spezies Reaktivität	Human
Immunogen	Recombinant full-length human protein was used as the immunogen for the TGM2 antibody.
Produktbeschreibung	Recognizes a 77-85kDa protein, identified as cellular or tissue transglutaminase II (TGase II, TGM2). Transglutaminases are enzymes that catalyze the crosslinking of proteins by epsilon-gamma glutamyl lysine isopeptide bonds. While the primary struct...
Klonalität	Monoclonal
Klon-Bezeichnung	[SPM592]
UniProt	P21980
Reinheit	Protein G affinity chromatography

Formulierung	Prediluted in 1X PBS, 0.1 mg/ml BSA (US sourced), 0.05% sodium azide, *For IHC use only*
Antibody Type	Primary Antibody
Application Verdünnung	The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.
Anwendungsbeschreibung	Optimal dilution of the TGM2 antibody should be determined by the researcher.1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min.2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.